arXiv:1412.1644v2 [math.CV] 20 Mar 2015

SHARP MARKOV-TYPE INEQUALITIES FOR RATIONAL FUNCTIONS ON SEVERAL INTERVALS

M. A. AKTURK AND A. LUKASHOV

Abstract. In this paper, we give sharp Rusak- and Markov-type inequalities for rational functions on several intervals when the system of intervals is a "rational function inverse image" of an interval and those functions are large in gaps.

1. Introduction

In 1889 A. A. Markov proved that if $P_n(x)$ is a polynomial of degree n and $|P_n(x)| \le M$ on an interval of length $2L$, then $|P_n'(x)| \le n^2 ML$ on the same interval and factor n^2 is attained for (transformed) Chebyshev polynomials of the first kind. Reader may consult papers [4, 20] and books [5, 18] for more detailed story of this inequality and its generalizations.

It is clear that Markov inequality can not be generalized for the set of all rational functions of degree n, although there are some Markov-type inequalities for rational functions with free poles [7, 8, 10]. For rational functions with fixed denominators under additional restrictions on the poles sharp Markov-type inequalities on an interval were obtained by V.N. Rusak [19].

Generalizations of Markov inequality onto polynomials and rational function for sets more complicated then one interval were considered in many papers [9, 16, 17, 21], but to the best of our knowledge the sharp factor for Markov inequalities is not known even for the simplest case of polynomials of even degree on two symmetric intervals. Asymptotically sharp Markov-type inequalities for polynomials on several intervals were obtained in [6, 22]. We start with one result of V.N. Rusak which was the base for sharp Markov type inequalities on one interval for rational functions with fixed poles.

Theorem 1.1. [19] *If algebraic fraction of the from*

$$r_n(x) = \frac{p_n(x)}{\sqrt{t_{2n}(x)}},$$

$$t_{2n}(x) = \prod_{k=1}^{2n} (1 + a_k x)\,, \ -1 \le x \le 1, \tag{1.1}$$

Date: July 11, 2014 and, in revised form, Month Day, Year.

2010 *Mathematics Subject Classification.* 41A17, 41A20.

Key words and phrases. Inequalities in approximation, Approximation by rational functions.

The authors were supported by Scientific and Technological Research Council of Turkey (TUBİTAK), joint project with Russian Foundation of Basic Research (RFBR), no.113F369.

where $p_n(x)$ is an algebraic polynomial of degree at most n with complex (real) coefficients, numbers a_k are either real or pairwise complex conjugate, $|a_k|<1, k=1,\dots,2n$, satisfies the condition

$$|r_n(x)|\le 1,\ -1\le x\le 1, \tag{1.2}$$

then the estimate

$$|r_n'(x)|\le \begin{cases} \frac{\lambda_n(x)}{\sqrt{1-x^2}},\ x_1\le x\le x_n, \\ |m_n'(x)|,\ -1\le x\le x_1,\ x_n\le x\le 1, \end{cases} \tag{1.3}$$

is valid. Here $\{x_k\}$, $-1<x_1<\ldots<x_n<1$ are zeros of the cosine fraction $m_n(x)=\cos\frac{1}{2}\sum\limits_{k=1}^{2n}\arccos\left(\frac{x+a_k}{1+a_kx}\right)$, and $\lambda_n(x)=\frac{1}{2}\sum\limits_{k=1}^{2n}\frac{\sqrt{1-a_k^2}}{1+a_kx}$. Equality sign in (1.3) is attained only for functions $r_n(x)\equiv\varepsilon m_n(x),\ |\varepsilon|=1$.

The main goal of the paper is to give maximal (in a sense) possible extension of Rusak's inequality for rational functions on several intervals. As a consequence we give sharp Markov-type inequality for rational functions with an additional restriction. Note that the equality for our Markov-type inequality is attained for rational functions which can be considered as an analogue of Chebyshev-Markov rational functions (their numerators are called also Bernstein-Szegő orthogonal polynomials) on several intervals (see, for example [12, 13, 14, 15]). We would like to remark that general result of V. Totik [22] contains $o(1)$ as $n\longrightarrow\infty$, hence it says nothing for a fixed (maybe small) n.

2. Results

Let $\Re(\xi_1,\dots,\xi_{2n})$ be the set of all " rational functions "of the form

$$r(x)=\frac{b_0x^n+b_1x^{n-1}+\ldots+b_n}{\sqrt{\rho_\nu(x)}}, \tag{2.1}$$

$b_0,\dots,b_n\in\mathbb{C}$ and $\rho_\nu(x)=\prod\limits_{j=1}^{2n}(x-\xi_j)$ is a real polynomial of degree ν which is positive on $E=\bigcup\limits_{j=1}^{l}[a_{2j-1},a_{2j}]$, $-1=a_1<a_2<\ldots<a_{2l}=1$ (ξ_j might be equal to ∞, then $(x-\xi_j)$ should be omitted) and

$$\varpi_E(z,x)=\frac{\partial}{\partial x}\omega\left(z,E\cap[a_1,x],\mathbb{C}\backslash E\right);\omega_k(\xi_j)=\omega\left(\xi_j,[a_{2k-1},a_{2k}],\mathbb{C}\backslash E\right),k=1,\dots,l.$$

Here $\omega(z,G,\mathbb{C}\backslash E)$ is the harmonic measure of a set $G\subset E$ at a point $z\in\mathbb{C}\backslash E$.

Consider also a subclass $\Re^*(\xi_1,\dots,\xi_{2n})$ of $\Re(\xi_1,\dots,\xi_{2n})$ which consists of those functions $r\in\Re(\xi_1,\dots,\xi_{2n})$, which satisfy $|r(x)|>\|r\|_{C(E)}$ for all $x\in[-1,1]\backslash E$.

Theorem 2.1. *Suppose that $\sum\limits_{j=1}^{2n}\omega_k(\xi_j)=2q_k,\ q_k\in\mathbb{N},\ k=1,\dots,l$, and $|\xi_j|>1, j=1,\dots,2n$. Then for any $r\in\Re^*(\xi_1,\dots,\xi_{2n}),\|r\|_{C(E)}=1$ the inequality*

$$|r'(x)|\le\begin{cases}\gamma_n'(x),x\in\widetilde{E}_n,\\ |m_n'(x)|,x\in E\backslash\widetilde{E}_n\end{cases} \tag{2.2}$$

is valid, where

$$\begin{aligned} m_n(x) &= \cos\left(\gamma_n\left(x\right)\right), \\ \gamma_n\left(x\right) &= \frac{\pi}{2}\int\limits_{a_1}^{x}\sum_{j=1}^{2n}\varpi_E\left(x,\xi_j\right)dx, \\ \widetilde{E}_n &= \left[x_1,x_{q_1}\right]\cup\left[x_{q_1},x_{q_1+q_2}\right]\cup\ldots\cup\left[x_{q_1+\ldots+q_{l-1}},x_n\right], \end{aligned}$$

and $x_1<\ldots<x_n$ *are zeros of* m_n *(there are* q_k *zeros on* $[a_{2k-1},a_{2k}]$, $k=1,\ldots,l$*).*

For $r(x)\equiv\varepsilon m_n(x)$, $|\varepsilon|=1$, *inequality in (2.2) is attained.*

Proof. It is sufficient (compare [5, proof of Corollary 5.1.5]) to prove Theorem 2.1 only for the case when numerator $p_n(x)$ has real coefficients. Firstly consider the case $x\in\widetilde{E}_n$. Then by [14, Theorem 5]

$$|r'\left(x\right)|\leq\sqrt{1-r^2\left(x\right)}\gamma_n'\left(x\right)\leq\gamma_n'\left(x\right) \tag{2.3}$$

Next we consider the case $x\in E\backslash\widetilde{E}_n$. Here we adopt Rusak's method [19, p. 58-60].

Let $\{y_j\}_{j=1}^{n+l}$ be zeros of the sine Chebyshev-Markov fraction $\frac{\sqrt{-H(x)}v_{n-l}(x)}{\sqrt{\rho_\nu(x)}}$, where $v_{n-l}(x)$ is determined from

$$m_n^2(x)-\frac{H(x)v_{n-l}^2(x)}{\rho_\nu(x)}=1,$$

and $H(x)=\prod\limits_{j=1}^{2l}\left(x-a_j\right)$ (see [14, Theorem 3]). Note that m_n is a function of the form (2.1) and

$$m_n(y_k) = (-1)^{\mu+j},\quad y_k\in[a_{2j-1},a_{2j}],\quad k=1,\ldots n+l \tag{2.4}$$

$$k = \sum_{i=1}^{j-1}q_i+\mu+j-1,\ 1\leq\mu\leq q_j+1,\ j=1,\ldots,l. \tag{2.5}$$

If $r(x)$ is different from $m_n(x)$ then for $0<\lambda<1$ the difference $m_n(x)-\lambda\ r(x)$ takes the same signs as m_n at points $\{y_j\}_{j=1}^{n+l}$. Hence it has simple zeros on the set E and can be written as

$$m_n(x)-\lambda r(x)=C\frac{\prod\limits_{k=1}^{n}\left(x-z_k\right)}{\sqrt{\prod\limits_{j=1}^{2n}\left(x-\xi_j\right)}}.$$

Next, the derivative of this difference is equal to

$$m_n'(x)-\lambda r'(x)=\frac{C}{2}\left(m_n(x)-\lambda r(x)\right)\left[2\sum_{k=1}^{n}\frac{1}{x-z_k}-\sum_{j=1}^{2n}\frac{1}{x-\xi_j}\right]. \tag{2.6}$$

Zeros of the difference $m_n'(x)-\lambda r'(x)$ coincide with zeros of square brackets in (2.6), and they are zeros of the logarithmic derivative of the rational function $R_{2n}(x)=\prod\limits_{k=1}^{n}\left(x-z_k\right)^2/\rho_\nu(x)$.

The condition $|\xi_j| > 1,\ j = 1, \dots, 2n$ guarantees that all poles of the rational function $R_{2n}(z)$ are contained in region $|z| \geq r > 1$, all zeros $\{z_k\}$ of the function $\{R_{2n}(z)\}$ lie in the interval $(-1, 1) \supset E$.

Bocher-Walsch Theorem [23, Corollary 1 on p. 99] implies that $R'_{2n}(z)$ has $n-1$ zeros in $|z| \leq 1$ and other zeros are in $|z| \geq r > 1$. In the same way because of (2.6) the zeros of the difference $m'_n(x) - \lambda r'(x)$ are located. So the function $m'_n(x) - \lambda r'(x),\ 0 < \lambda < 1$ has no more than $n-1$ zeros in $(-1, 1)$ and other zeros are in $|z| \geq r > 1$.

By the continuity as λ tends to 1 we obtain the function $m'_n(x) - r'(x)$ has at most $n-1$ zeros on $[-1, 1]$.

From the identity

$$\left(\frac{m'_n(x)}{\gamma'_n(x)}\right)^2 + m_n^2(x) = 1,$$

and the inequality (with $\leq$sign it was given in [14, Theorem 5], for $r \neq \pm m_n$ strict inequality was obtained in [11])

$$\frac{(r'(x))^2}{\gamma'_n(x)} + r^2(x) < 1,$$

we get

$$|r'(x_k)| < |m'_n(x_k)|.$$

Since

$$m'_n(x) = -\sin\gamma_{2n}(x)\,\gamma'_n(x),$$

has alternate sign at points $\{x_k\}_{k=1}^n$, the relation $sign\,\{m'_n(x_k) - r'(x_k)\} = (-1)^k$, $k = 1, \dots, n$ holds. On the interval (x_1, x_{q_1}) there are $q_1 - 1$ zeros of $m'_n(x) - r'(x)$, on $(x_{q_1+1}, x_{q_1+q_2})$ there are $q_2 - 1$ zeros, and so on, therefore $n - l$ zeros of the difference $m'_n(x) - r'(x)$ are on the set $\widetilde{E}_n$. Since $m_n(x) - \lambda r(x)$ has the same value at a_{2k} and $a_{2k+1}, k = 1, \dots, l-1$, there is at least one zero of $m'_n(x) - \lambda r'(x)$ on $(a_{2k}, a_{2k+1}), k = 1, \dots, l-1$. So by the continuity as λ tends to 1 we obtain that $m'_n(x) - r'(x)$ preserves sign inside intervals from $E \backslash \widetilde{E}_n$.

In particular it follows from $m'_n(a_1) - r'(a_1) \leq 0$ that $m'_n(x) - r'(x) \leq 0$ for $a_1 \leq x \leq x_1$. Analogous consideration is valid for the algebraic fraction $-r(x)$. Therefore we obtain inequality $m'_n(x) + r'(x) \leq 0$ for $a_1 \leq x \leq x_1$, and it gives the estimate $|r'(x)| \leq |m'_n(x)|,\ a_1 \leq x \leq x_1$. Analogously other intervals from $E \backslash \widetilde{E}_n$ are considered.

□

Theorem 2.2. *Suppose that* $\sum_{j=1}^{2n} \omega_k(\xi_j) = 2q_k,\ q_k \in \mathbb{N},\ k = 1, \dots, l,$ *and* $\xi_j \in \overline{\mathbb{R}}, |\xi_j| > 1, j = 1, \dots, 2n$. *Then for any* $r \in \Re^*(\xi_1, \dots, \xi_{2n}), \|r\|_{C(E)} = 1$ *the inequality*

$$\|r'\|_{C(E)} \leq \|m'_n\|_{C(E)} \tag{2.7}$$

holds.

Proof. Firstly consider the case $x \in \widetilde{E}_n$. Then by Theorem 2.1

$$\begin{aligned} |r'(x)| &\leq \gamma'_n(x) \\ &\leq \max_{x \in \widetilde{E}_n} \gamma'_n(x). \end{aligned}$$

By [2] $\gamma_n'(x)$ is convex hence

$$\max_{x\in\widetilde{E}_n}\left(\gamma_n'(x)\right)=\max_{k=1,\dots,n}\left(\gamma_n'(x_k)\right)$$

and we get by

$$\gamma_n'(x_k)=\left|m_n'(x_k)\right|, k=1,\dots,n,$$

the desired result,

$$|r'(x)|\le\|m_n'\|_{C(E)}, x\in\widetilde{E}_n,$$

and application of second inequality in Theorem 2.1 finishes the proof.

□

Corollary 2.3. Let Π_n denote the set of algebraic polynomials with real coefficients of degree at most n. If $0<a<b$, n is even and $p_n\in\Pi_n^*$ where Π_n^* is the subclass of Π_n which consists of polynomials $p_n\in\Pi_n$ such that $p_n(x)>\|p_n\|_{[-b,-a]\cup[a,b]}$ for $x\in(-a,a)$, then the inequality

$$\|p_n'\|_{[-b,-a]\cup[a,b]}\le\frac{n^2b}{b^2-a^2}\|p_n\|_{[-b,-a]\cup[a,b]}.$$

is valid. It is attained for polynomials

$$S_n(x)=T_{n/2}\left(\frac{2x^2-b^2-a^2}{b^2-a^2}\right),$$

where T_n is the n^{th} Chebyshev polynomial $(T_n=\cos(n\arccos(x)))$.

Proof. It is sufficient to note that for $E=[-b,-a]\cup[a,b]$.

$$\varpi_E(x,\infty)=\frac{1}{\pi}\frac{|x|}{\sqrt{(b^2-x^2)(x^2-a^2)}},$$

and for $\xi_1=\ldots=\xi_{2n}=\infty, n$ is even, $m_n(x)=S_n(x)$.

□

Remark 2.4. Corollary 2.3 complements [6, Theorem 2].

Remark 2.5. The condition $r\in\Re^*(\xi_1,\dots,\xi_{2n})$ is essential for validity of Theorem 2.1. Indeed, let $E=[-1,-a]\cup[a,1], \rho_\nu(x)=1$. Here

$$m_4(x)=\frac{8x^4-8x^2(1+a^2)+1+6a^2+a^4}{(1-a^2)^2}$$

and

$$m_4'(a)=-\frac{16a}{1-a^2}.$$

But for the Chebyshev polynomial $T_3(x)=4x^3-3x$ we have

$$T_3'(a)=12a^2-3,$$

hence

$$|T_3'(a)|>|m_4'(a)|$$

for sufficiently small a. We don't know if the inequality in Theorem 2.2 is valid for $r\in\Re(\xi_1,\dots,\xi_{2n})\setminus\Re^*(\xi_1,\dots,\xi_{2n})$ or not.

Remark 2.6. The main result of [1] which corresponds to Theorem 2.1, is valid under additional supposition $|r(x)|>\|r\|_{C(E)}$ for all $x\in conv(E)\backslash E$.

Remark 2.7. The supposition ($|\xi_j| > 1$) is essential even for the case of one interval, as the example (taken from [19]) $r(x) = \frac{72x^2-5x}{100x^2+1}$ shows (here $\|r'\|_{[-1,1]} \geq r'(1) > \|m_2'\|_{[-1,1]}, m_2(x) = \frac{102x^2-1}{100x^2+1}, \|r\|_{[-1,1]} \leq 1$).

ACKNOWLEDGEMENTS. The authors are deeply grateful to reviewers and to Professor V. Totik for careful reading of the manuscript and remarks which helped to improve the text and correct mistakes.

References

1. M. A. Akturk, and A. Lukashov, *Markov-type inequalities for rational functions on several intervals*, AIP Conf. Proc. **1611** (2014), 208-210.
2. D. Benko, P. Dragnev, and V. Totik, *Convexity of harmonic densities*, Rev. Mat. Iberoam. **28(4)** (2012), 1-14.
3. D. Benko, and V. Totik, *Sets with Interior Extremal Points for the Markoff Inequality*, Journal of Approximation Theory **110** (2001), 261–265.
4. R. P. Boas, *Extremal Problems for Polynomials*, The American Mathematical Monthly, **85(6)** (1978), 473–475.
5. P. Borwein, and T. Erdélyi, *Polynomials and Polynomial Inequalities*, Springer, New York, 1995.
6. P. Borwein, *Markov's and Bernstein's inequalities on disjoint intervals*, Canad. Math.J. **38(1)** (1981), 201–209.
7. V. I. Danchenko, *Several integral estimates of the derivatives of rational functions on sets of finite density*, Mat. Sb. **187(10)** (1996), 33-52; English transl., Sb. Math. **187** (1996), 1443-1463.
8. E. P. Dolzhenko, *Estimates for derivatives of rational functions*, Izv. Akad. Nauk SSSR Ser. Mat. **27(1)** (1963), 9-28. (Russian)
9. T. Erdélyi, A. Kroó, and J. Szabados, *Markov-Bernstein type inequalities on compact subsets of R*, Analysis Mathematica, **26** (2000), 17-34.
10. A. A. Gonchar, *Inverse theorems on the best approximation by rational functions*, Izv. Akad. Nauk SSSR Ser. Mat. **25(2)** (1961), 347-356. (Russian)
11. S. I. Kalmykov, *Majoration Principles and some Inequalities for Polynomials and Rational Functions with Prescribed Poles*, J. Math. Sci. **157(4)** (2009), 623–631.
12. V. I. Lebedev, *Extremal Chebyshev-Markov-Bernstein-Szegő Polynomial of the 1-4 kind and methods for the optimization of computational algorithms*, Numerical methods for solving linear and nonlinear boundary value problems(Russian), 37–87, Tr. Mat. Tsentra im. N. I. Lobachevskogo, **20** Kazan. Mat. Obs., Kazan (2003).
13. A. L. Lukashov, *On Chebyshev-Markov Rational Functions over Several Intervals*, Journal of Approximation Theory, **95** (1998), 333–352.
14. A. L. Lukashov, *Inequalities for Derivatives of Rational Functions on Several Intervals*, Izv. Math. **68(3)** (2004), 543–565.
15. F. Peherstorfer, *Orthogonal and extremal polynomials on several intervals*, Journal of Computational and Applied Mathematics, **48** (1993), 187–205.
16. W. Pleśniak, *Inégalité de Markov en Plusieurs Variables*, International Journal of Mathematics and Mathematical Sciences, (2006), Article ID 24549, 1-12.
17. A. A. Privalov, *An analogue of A. A. Markov's inequality, Application to interpolation and Fourier Series*, Orthogonal series and approximation of functions, **164** (1983), 142–154. (Russian)
18. Q.I. Rahman, and G. Schmeisser, *Analytic Theory of Polynomials*, Clarendon Press, Oxford, 2002.
19. V. N. Rusak, *Rational Functions as an Apparatus of Approximation*, Belorus. State Univ. Press, Minsk, 1979. (in Russian)
20. A. Shadrin, *Twelve proofs of the Markov inequality*, in: Approximation Theory: a volume dedicated to Borislav Bojanov, Prof. Drinov Acad. Publ. House, Sofia, (2004), 233–298.
21. V. Totik, *On Markoffs Inequality*, Constr. Approx., **18** (2002), 427-441.
22. V. Totik, *Polynomial Inverse Images and Polynomial Inequalities*, Acta Math. **187(1)** (2001), 139–160.

23. J.L. Walsh, *The Location of Critical Points of Analytic and Harmonic Functions*, AMS, New York, 1950.

Department of Engineering Sciences, Istanbul University, 34320 Istanbul, Turkey
Current address: Department of Engineering Sciences, Istanbul University, 34320 Istanbul, Turkey
E-mail address: mehmetaliakturk@yandex.com

Department of Mathematics, Fatih University, 34500 Istanbul, Turkey, and Department of Mechanics and Mathematics, Saratov State University, 410012 Saratov, Russia
E-mail address: alukashov@fatih.edu.tr